

Updated 24 April 2020

Frequently Asked Questions for KCDC on COVID-19

(Updated on 24 April 2020)

1. Diagnostic Testing

Q1. Who can get tested for COVID-19? Do you test asymptomatic persons too?

We test persons who are "suspected cases" or "Patients Under Investigation (PUI)" as defined by our COVID-19 Response Guidelines (excerpt provided below).

Excerpt from COVID-19 Response Guidelines (Edition 7-4):

1. Case Definition

• Suspected case:

A person exhibiting fever (37.5 degrees or above) or respiratory symptoms (coughs, shortness of breath, etc.) within 14 days of contact with a confirmed COVID-19 patient during the confirmed patient's symptom-exhibiting period.

• Patient Under Investigation (PUI):

(1) A person suspected of COVID-19 according to a physician's opinion for reasons such as pneumonia of an unknown cause;

(2) A person exhibiting fever (37.5 degrees or above) or respiratory symptoms (coughs, shortness of breath, etc.) within 14 days of entering Korea after visiting another country; or

(3) A person exhibiting fever (37.5 degrees or above) or respiratory symptoms (coughs, shortness of breath, etc.) with an epidemiological link to a domestic COVID-19 cluster.

We do sometimes apply exceptions for certain high risk groups. For example, we tested all persons linked to certain major clusters (i.e., Shincheonji, Guro-gu call center) regardless of clinical symptoms. We have also recently tested all persons in long-term care facilities in Daegu City regardless of clinical symptoms.

Q2. What kind of diagnostic tests do you use for COVID-19? How is the test performed?

Korea uses COVID-19 nucleic acid testing (real-time RT-PCR) to diagnose a patient.

Specimens are collected by doctors, nurses, or clinical laboratory scientists at designated locations (i.e., screening centers/stations). Upper respiratory tract specimens are required. Lower respiratory tract specimens are also collected if the patient has sputum. Nucleic acid testing may be performed directly at screening centers if the center is equipped with such capacity; otherwise, the specimens are sent to a testing center.

Excerpt from COVID-19 Response Guidelines: Specimen Collection Method

• Upper respiratory tract specimen:

Nasopharyngeal and oropharyngeal swab mixture (1 tube)



Updated 24 April 2020

- (For nasopharyngeal swab) Insert a swab deep into nostril.

- (For oropharyngeal swab) Scrape from the inner wall of the throat with a swab.

• Lower respiratory tract specimen:

Have patient cough deeply and collect the sputum. Ensure that saliva is not included.

If patient does not have sputum, do not induce sputum as spitting forcibly may cause aerosol.

Q3. What is your current diagnostic testing capacity? How was Korea able to expand testing capacity in a short period of time?

Currently, there are a total of 118 institutions available for diagnostic tests: Korea Centers for Disease Control and Prevention (1), National Quarantine Stations (4), Institutes of Health and Environment (18), private clinical laboratories and hospitals (95). On average, 15,000 tests (maximum 20,000) can be performed per day.

The relatively fast expansion of testing capacity was made possible thanks to active collaborative efforts between government, academia, and private sector.

Korean government (KCDC) quickly developed a test and disclosed it. Based on this, a company developed and produced diagnostic reagent. Upon evaluation by government and medical academic experts, the reagent was granted emergency use authorization by the Ministry of Food and Drug Safety. Testing facilities across the nation then began using the test.

To ensure accuracy of the tests performed, COVID-19 testing centers were selected by KCDC from testing facilities that had been certified for outstanding quality. They also received additional training and passed external quality assessment. Testing quality of each center is maintained by quality assurance by KCDC and academic experts.

Q4. Do you conduct pooled testing for COVID-19?

To improve testing efficiency, KCDC and Korean Society for Laboratory Medicine have jointly prepared a protocol for pooled testing, which is a method in which biological samples from multiple people are combined into a testing pool and tested via a single test. In the event of positive result, individual testing is carried out using previously reserved samples. This method is useful for screening asymptomatic high-risk groups. We are starting pooled testing for certain high-risk facilities, such as long-term care facilities, with exposure risk high enough to warrant having entire staff and patients.

2. Epidemiological Investigation

Q1. What methods of contact tracing do you use for COVID-19?

Contact tracing is largely divided into 4 stages: (1) investigation, (2) exposure risk assessment, (3) contact classification, and (4) contact management. During the investigation phase, basic



Updated 24 April 2020

information including whereabouts of the patient for a certain period of time is collected through the process of interviewing patients. Family or healthcare workers may also be interviewed if needed. If supplementary information is needed (e.g., due to memory omission or inconsistencies), more objective information (such as medical records, cellular GPS data, credit card transactions, CCTV footages) may be collected during the risk assessment stage. Contacts identified based on the collected information are subject to self-quarantine (home quarantine) along with health education and symptom monitoring.

Q2. What is the basis on which you collect and/or use personal data in epidemiological investigation?

Information required for epidemiological investigation can be collected and/or used within the scope permitted by the Infectious Disease Control and Prevention Act.

Excerpt from Infectious Disease Control and Prevention Act:

Article 76-2 (Request to Provide Information, etc.)

(1) If necessary to prevent infectious diseases and block the spread of infection, the Minister of Health and Welfare or the Director of the Korea Centers for Disease Control and Prevention may request the heads of relevant central administrative agencies (including affiliated agencies and responsible administrative agencies thereof), the heads of local governments (including superintendents of education prescribed in Article 18 of the Local Education Autonomy Act), public institutions designated under Article 4 of the Act on the Management of Public Institutions, medical institutions, pharmacies, corporations, organizations, and individuals to provide the following information concerning patients, etc. with infectious diseases and persons likely to be infected by infectious diseases, and persons in receipt of such request shall comply therewith: <Amended by Act No. 14286, Dec. 2, 2016>

1. Personal information, such as names, resident registration numbers prescribed in Article 7-2 (1) of the Resident Registration Act, addresses, and telephone numbers (including cell phone numbers);

2. Prescriptions prescribed in Article 17 of the Medical Service Act, records of medical treatment prescribed in Article 22 of the same Act, etc.;

3. Records of immigration control during the period determined by the Minister of Health and Welfare;

4. Other information prescribed by Presidential Decree for monitoring the movement paths of patients with infectious diseases.

(2) If necessary to prevent infectious diseases and block the spread of infection, the Minister of Health and Welfare may request the relevant head of the National Police Agency, regional police agency, and police station established under Article 2 of the Police Act (hereafter in this Article, referred to as "police agency") to provide location information of patients, etc. with an infectious disease and persons likely to be infected by an infectious disease. In such cases, notwithstanding Article 15 of the Act on the Protection, Use, etc. of Location Information and Article 3 of the Protection of Communications Secrets Act, the relevant head of a police agency, upon request by the Minister of Health and Welfare, may request any location information provider defined in Article 5 (7) of the Act on the Protection, Use, etc. of Location Information susiness operator defined in subparagraph 8 of Article 2 of the Telecommunications Business Act, to provide location information of patients, etc. with an infectious disease and persons likely to be infected by an infectious disease; and the location information provider and the telecommunications business operator in receipt of such request shall comply therewith, except in extenuating circumstances. <Amended by Act No. 13639, Dec. 29, 2015>



Updated 24 April 2020

(3) The Minister of Health and Welfare may provide information collected pursuant to paragraphs (1) and (2) to the heads of the relevant central administrative agencies, the heads of local governments, the chairperson of the National Health Insurance Corporation, the president of the Health Insurance Review and Assessment Service, and such medical personnel, medical institutions, and other organizations as are performing tasks related to infectious diseases. In such cases, information provided shall be limited to information related to the tasks of the relevant agencies, etc., for preventing infectious diseases and blocking the spread of infection.

Q3. How do you deal with privacy issues regarding collection and use of personal data for epidemiological investigation?

Privacy-sensitive data such as cellular GPS logs, credit card transaction logs, and camera footages are NOT collected for every confirmed COVID-19 patient. The need to collect such data is carefully assessed based on the result of preliminary investigation through patient interviews. From our experience, we have found that some patients provide inaccurate or partial information during interview, either unintentionally or deliberately, while some others have trouble recalling every location they have visited from their memory. For certain infectious diseases, such information gaps may be detrimental to disease control efforts and put the general population at great risk. Our investigators do their best to gather as much information as possible by first interviewing patients, their family, and/or healthcare workers. If, despite their efforts, there is still not enough information to identify the infection source and/or contacts, relevant personal data is collected if available. Information collected during epidemiological investigation is managed as safely as possible with strict security measures in place and is destroyed when the investigation is over, as required by law.

Q4. How much contact tracing / epidemiological investigation capacity does Korea have in terms of human resources?

Since the MERS outbreak in 2015, there has been effort to regularly recruit and train epidemiological intelligence (EI) officers at both central and local levels. Currently, there are 75 EI officers at central level and 54 at city or province level. We also have a pool of around 300 private epidemiologists that have been appointed to work on COVID-19 response. Local public health centers (around 256 nationwide) also have capacity to conduct basic contact tracing and preliminary epidemiological investigation.

Korea also has 11 local-level support centers for control and prevention of infectious diseases. Although these centers are not governmental organizations, their management, operation, and allocation of budget is based on private-public partnership. In normal times, they work on training of EI officers, surveillance on infectious diseases, and providing advices on control and prevention of infectious diseases. During emergency, they can take part in as a response team. For example, during early stages of COVID-19 outbreak, KCDC's rapid response team was sent to the areas where confirmed cases were reported and conducted all epidemiological



Updated 24 April 2020

investigation. However, as the number of confirmed cases soared, these support centers took over the role of conducting preliminary investigation within their city or province, working in partnership with local public health authorities. Rapid response teams are sent out only when necessary (i.e. when further, in-depth investigation is required or when big clusters occur). This collaborative partnership greatly eased the resource strain on central government and its staff and enabled more efficient utilization of human resources.

3. Case Management

Q1. Korea has a relatively low fatality rate for COVID-19 cases. What special strategies do you believe contributed to achieving this?

As the number of patients rapidly surged in Daegu City and Gyeongbuk Province starting late February (which caused shortage in healthcare resources), we created a new system for allocating hospital beds based on the severity of the patient. This system enabled more medical resources to go to the more severe patients in need of urgent care, allowing more efficient treatment and management of patients.

In addition, persons who are considered high-risk groups are classified as severe patients regardless of clinical symptoms, so that they can receive timely care if needed.

COVID-19 high-risk groups are:

- The elderly 65 years or older;

- Persons with underlying chronic conditions, such as diabetes, chronic renal, liver, lung or cardiovascular diseases, or HIV;
- Patients with a blood cancer;
- Cancer patients receiving chemotherapy;
- Patients taking one or more immunosuppressant drugs;
- Pregnant women, patients with extreme obesity, dialysis patients, transplant patients, or smokers; or
- Patients with blood oxygen saturation level below 90%

Q2. Are all confirmed COVID-19 patients admitted to and treated at hospitals?

As mentioned above, patients confirmed with COVID-19 are classified into 4 categories based on severity: Mild, Moderate, Severe, and Extremely Severe. Mild patients (a.k.a., patients with mild symptoms) are placed in home quarantine (self-quarantine) or admitted to a Living Treatment Center and may later be transferred to a hospital if their symptoms aggravate. Moderate patients are admitted to hospitals with capacity to treat infectious disease patients. Severe or Extremely Severe patients are admitted to tertiary hospitals which have greater capacity for treatment. This system allows patients to receive the care they need in a timely manner.



Updated 24 April 2020

Q3. Any statistical information on patients who tested positive again after being discharged from isolation? (sometimes also referred to as "retested-positive cases" or "re-positive cases")

As of 17 April, of all 7,829 cases that have been discharged from isolation, 163 (2.1%) have tested positive again. On average, it took 13.5 days (min. 1 day – max. 35 days) between discharge from isolation and retesting positive. Out of 137 cases whose clinical and epidemiological information have been identified, 61 were symptomatic; 72 were asymptomatic; and 4 were under investigation. All of the 61 symptomatic cases had mild symptoms. No secondary infection from the retested-positive cases has been reported yet.

Further demographic information of these re-tested positive cases is as indicated in the table below.

		n	(%)
Total (as of 0:00 of 17 April)		163	(100.0%)
Sex	Male	54	(33.1%)
	Female	109	(66.9%)
Age group	80 and above	15	(9.2%)
	70-79	6	(3.7%)
	60-69	18	(11.0%)
	50-59	32	(19.6%)
	40-49	19	(11.7%)
	30-39	24	(14.7%)
	20-29	38	(23.3%)
	10-19	6	(3.7%)
	10 and below	5	(3.1%)
Location reported	Seoul	7	(4.3%)
	Daegu	67	(41.1%)
	Incheon	1	(0.6%)
	Daejeon	1	(0.6%)
	Ulsan	2	(1.2%)
	Sejong	5	(3.1%)
	Gyeonggi	13	(8.0%)
	Gangwon	3	(1.8%)
	Chungbuk	2	(1.2%)
	Chungnam	3	(1.8%)
	Gyeongbuk	54	(33.1%)
	Gyeongnam	3	(1.8%)
	Jeju	1	(0.6%)
	Point of Entry	1	(0.6%)

* More information will be available once the on-going investigation, research, and case studies are completed.