

දුරකථන) 0112669192 , 0112675011
தொலைபேசி) 0112698507 , 0112694033
Telephone) 0112675449 , 0112675280

ෆැක්ස්) 0112693866
பெக்ஸ்) 0112693869
Fax) 0112692913

විද්‍යුත් තැපෑල) postmaster@health.gov.lk
மின்னஞ்சல் முகவரி)
e-mail)

වෙබ් අඩවිය) www.health.gov.lk
இணையத்தளம்)
website)



සුවසිරිපාය
சுவசிரிபாய
SUWASIRIPAYA

සෞඛ්‍ය අමාත්‍යාංශය
சுகாதார அமைச்சு
Ministry of Health

මගේ අංකය) EPID/400/n-CoV/Vaccine_22
எனது இல)
My No.)

ඔබේ අංකය)
உமது இல)
Your No.)

දිනය)
திகதி)
Date)

2022/5/5

DDG- NHSL, NH Kandy
Directors of all hospitals,
Provincial Directors of Health Services
Regional Directors of Health Service

Second booster dose of Covid 19 vaccine: COMIRNATY - Covid 19 mRNA (Nucleoside modified -Pfizer BNT) vaccination campaign, from 05/05//2022

- 1. All aged 60 years of age and above, at least 3 months after the 1st booster dose**
- 2. Any one from 20 to 59 years: on request, at least 3 months after the 1st booster dose**

The Ministry of Health has taken the decision to provide a 2nd booster vaccination essentially for the category aged 60 years and above and anyone request for the age group 20 to 59 years, keeping the duration of at least 3 months after the 1st booster vaccination dose, from 05/05/2022.

- The vaccine for the 2nd booster dose campaign will be the “COMIRNATY” - Covid 19 mRNA Vaccine (Nucleoside modified -Pfizer BNT).
- It is advised to provide a dose for eligible categories those who have completed 2 primary doses from any type of vaccine and one Pfizer BNT booster dose, and the time elapsed 3 month from the 1st booster vaccine dose (last vaccination dose).
- Advise to identify and vaccinate all 60 years and above age group in all MOH areas, medical clinic attendees and OPD attendees.
 - Deploy all public health staff to identify eligible 60 years and above category on house-to-house identification to ensure all will get the 2nd booster vaccination dose.
 - At the same time, if any defaulters for the 1st booster dose or for primary doses, need to be provide the missed dose at the earliest possible.
- Advise strictly to follow the COMIRNATY, Covid 19 mRNA (Nucleoside modified -Pfizer BNT) vaccination guidelines issued on 05/07/2021 by the Epidemiology Unit, Ministry of Health find in www.epid.gov.lk for vaccine storage, vial preparation, administration, prepare, follow up and surveillance of Adverse Events Following Immunization (AEFI). (Only deviation is administration of a single dose.
- Advise to use the same consent form for adult Covid-19 vaccination.
- Each vial of COMIRNATY, Covid 19 mRNA (Nucleoside modified -Pfizer BNT) has 6 doses to vaccinate eligible individuals (follow the guidelines issued on 05/07/2021 by the Epidemiology Unit, Ministry of Health).
- It is advised to manage vaccine stocks with the plan to get down additional stocks as with the requirement ensure all required categories provide the vaccination on their first contact.
- The vaccines taken out from the Ultra-Low Temperature (ULT) freezer (from -70⁰C) will not be able to re-freeze in ULT freezer.

- **Vaccines supplied to the RMSD, MOH offices and hospitals should be stored only in the 2^o-8^oC in the Ice Lined Refrigerator (ILR), adhering to the national vaccine storage guidelines issued by the Epidemiology Unit.**

- **Even though ULT freezer (-70^oC) facilities available in anywhere in any districts, Comirnaty-Pfizer vaccine issued after the Epidemiology Unit cold stores should not be stored at ULT freezer (-70^oC) as it is transported in 2^o-8^oC.**

- Need to maintain regular communications with the Consultant Epidemiologists at the Epidemiology Unit, Provincial and District Consultant Community Physicians and Regional Epidemiologists in respective districts on the vaccination process to receive required vaccine supply and technical guidance to ensure proper storage and achieving maximum coverage for booster doses.

- MOOH and hospital directors/ responsible focal persons in the hospital are responsible for the vaccine stock management to ensure proper supply of vaccines:
 - All COMIRNATY, Covid 19 mRNA (Nucleoside modified -Pfizer BNT) vaccine stocks receiving to the MOH office or hospital **need to be stored in the ILR/refrigerator, marked clearly and stored separately according to received date, as separate supplied stocks will be expired in 30 days after taking out from the ULT (even though the vaccine vial expiry date is a different future date).**
 - Need to closely monitor until complete the vaccination from initial supply of vaccines, on “1st- received-1st – use” basis to ensure all received vaccines will be used within 28 days from the receipt date (at MOH and at hospitals).
 - Should not keep vaccines until 28 days without utilized and in 14 days of the receipt need to identify whether there are any excess vaccine doses remaining. In such situations, required to communicate with the Regional Epidemiologist in the district and with Consultant Epidemiologists at the Epidemiology Unit immediately for required redistributions without delay.
 - If any remaining unopened vials are returned after the clinic session, need to store it separately in a “return unopen vaccine vial container”, separately labelled and stored in the ILR and need to first use returned unopened vials first on the next vaccination day.

- Regional Epidemiologists are advised to closely monitor,
 - COMIRNATY, Covid 19 mRNA (Nucleoside modified -Pfizer BNT) vaccine stocks at the RMSD and should not exceed >30 days on receiving vaccines supplied at one lot.
 - On receiving vaccines at different dates, the date received needs to be clearly labelled, stocks on separate dates need to be separately stored.
 - Option of re-distribution of vaccines within institutions needs to be considered at all instances before re-order vaccines from the main stores of the Epidemiology Unit.

- Advise strictly to adhere to all the precautions and vaccination principles in conducting vaccination clinics.
 - Eligibility screening needs to be done by a Medical Officer.

- Emergency tray with all essential medicines is advised to be ready at all times and advised to check at every morning according to the guidelines given by the Epidemiology Unit.
- Transport arrangements to nearest hospital needs to be planned and ready to be used in any emergency AEFI situation after the initial management.
- Adhere to all infection prevention precautions to ensure protection from Covid-19 during vaccination sessions.

Need to deploy adequate trained and competent staff for vaccination (vaccine vial preparation and vaccination): refer to the poster attached for relevant steps of vaccine vial preparation (Annexure 1) and videos available at the google drive (<https://drive.google.com/drive/folders/1kvudLAUbxNKv3Gmt5mnjkckibrJFXM5h?usp=sharing>) shared through emails.

Identify vaccination data entry teams to deploy at the vaccination centre, MOH office and hospitals to enter vaccination data in to the vaccine tracker.

At the end of each day, total number vaccinated needs to be provided to the Epidemiology Unit, copying to Regional Epidemiologist in respective districts.

If you need any further clarification, please contact the Chief Epidemiologist of the Epidemiology Unit. Additional resource materials can find on the website www.epid.gov.lk.

Thank you,



Dr. Asela Gunawardena
Director General of Health Services

Dr. ASELA GUNAWARDENA
Director General of Health Services
Ministry of Health
"Suwasiripaya"
385, Rev. Baddegama Wimalawansa Thero Mawatha,
Colombo 10.

Cc:

- i. Hon. Minister of Health
- ii. Secretary, Ministry of Health
- iii. Additional Secretary (Public Health Services)
- iv. Deputy Director General (PHS) -1
- v. Chief Epidemiologist
- vi. Director/MCH/FHB
- vii. Provincial and district CCPs
- viii. Regional Epidemiologists
- ix. MOO/MCH